

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment

Charts updated 17 January 2022

Page 1 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.
No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

COVID-19 Antiviral Therapies (Licensed and Under Clinical Investigation)

Please refer to the additional notes on the following pages for further details and complete dose recommendations.

Drug	Hepatic Impairment			References
	Mild (Childs Pugh A)	Moderate (Childs Pugh B)	Severe (Childs Pugh C)	
Azithromycin	100%	100%	Not recommended	1, 2
Bamlanivimab/ Etesevimab	100%	100% (no data available)	100% (no data available)	3
Casirivimab/ Imdevimab	100%	100% (limited data available)	100% (no data available)	4, 5
Chloroquine	100%	100%	100%	6, 7
Favipiravir	Consider dose adjustment. 1200 mg twice daily then 800 mg twice daily to day 5	Consider dose adjustment. 1200 mg twice daily then 800 mg twice daily to day 5	Consider dose adjustment. 800 mg twice daily then 400 mg twice daily to days 2-3	8, 9
Hydroxychloroquine	100%	100%, with caution and frequent ALT monitoring	Caution. Consider 50% (to a maximum of 400 mg) with frequent ALT monitoring	10, 11
Interferon beta	100%	Not recommended	Not recommended	12-15
Ivermectin	100%. Not studied in hepatic disease.	100%. Not studied in hepatic disease.	100%, with caution. Not studied in hepatic disease.	16
Molnupiravir	100%	100%	100%	17
Niclosamide	100%	100%	100%	18
Nirmatrelvir + ritonavir	100%	100%	Contraindicated	19, 20
Nitazoxanide	100%, with caution. Not studied in hepatic disease.	100%, with caution. Not studied in hepatic disease.	100%, with caution. Not studied in hepatic disease.	21
Remdesivir	100% (no data available)	100% (no data available)	100% (no data available)	22, 23
Ribavirin	100%	100%, with caution and close monitoring of renal function, FBC and LFTs with dose reduction if required.	100%, with caution and close monitoring of renal function, FBC and LFTs with dose reduction if required.	24, 25
Sotrovimab	100% (no data available)	100% (no data available)	100% (no data available)	26, 27
Tixagevimab/ Cilgavimab	100% (no data available)	100% (no data available)	100% (no data available)	28

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment

Charts updated 17 January 2022

Page 2 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister. No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

Notes - Antiviral Therapies (Licensed and Under Clinical Investigation)

Azithromycin	Hepatic elimination: use with caution in all degrees of hepatic impairment. Discontinue if signs of hepatic dysfunction.
Bamlanivimab/ Etesevimab	No dosage adjustment is recommended in patients with mild hepatic impairment. Based on population PK analysis, there is no difference in PK of bamlanivimab or etesevimab in patients with mild hepatic impairment compared to patients with normal hepatic function. Bamlanivimab and etesevimab have not been studied in patients with moderate or severe hepatic impairment.
Casirivimab/ Imdevimab	Casirivimab and imdevimab are not expected to undergo significant hepatic elimination. No dosage adjustment is required in individuals with mild hepatic impairment. The effect of hepatic impairment on the exposure of casirivimab and imdevimab was evaluated by population PK analysis in patients with mild hepatic impairment (n=586 for casirivimab; n=599 for imdevimab) (total bilirubin >1.0-1.5 x ULN and any aspartate aminotransferase). No clinically important differences in the exposure of casirivimab and imdevimab were found between patients with mild hepatic impairment and patients with normal hepatic function. Limited data (n=11) are available in patients with moderate hepatic impairment; no data are available in patients with severe hepatic impairment.
Chloroquine	Use with caution; monitor liver function tests and watch for toxicities.
Favipiravir	Could consider extending treatment duration in COVID as per duration for ongoing trials. Dosing as per study US109.
Hydroxychloroquine	Maximum dosage based on minimal data and risk of hepatotoxicity.
Interferon beta	Caution if ALT >2.5x ULN. Dose reduction advised if ALT >5x ULN. Discontinue if jaundice or clinical symptoms of liver disease.
Ivermectin	Ivermectin has not been studied in patients with hepatic impairment. Patients with abnormal liver tests prior to starting ivermectin should be carefully evaluated.
Molnupiravir	No dose adjustment is required for patients with hepatic impairment. The pharmacokinetics of molnupiravir and NHC has not been evaluated in patients with hepatic impairment. Preclinical data indicate that hepatic elimination is not expected to be a major route of NHC elimination therefore hepatic impairment is unlikely to affect NHC exposure.

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment

Charts updated 17 January 2022

Page 3 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.
No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

Notes - Antiviral Therapies (Licensed and Under Clinical Investigation) [continued]

Niclosamide	Niclosamide may be given safely to patients with liver diseases.
Nirmatrelvir + ritonavir	<p>Nirmatrelvir and ritonavir exposures were similar in subjects with normal hepatic function and those with moderate hepatic impairment. No dosage adjustment is needed for patients with either mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment.</p> <p>Nirmatrelvir/ritonavir is contraindicated in patients with severe hepatic impairment (Child-Pugh Class C) as it has not been studied and there are no pharmacokinetic or safety data available regarding its use in subjects with severe hepatic impairment.</p>
Nitazoxanide	The pharmacokinetics of nitazoxanide in patients with compromised hepatic function have not been studied.
Remdesivir	<p>The pharmacokinetics of remdesivir have not been evaluated in patients with hepatic impairment. The role of the liver in the metabolism of remdesivir is unknown. It is not known if dosage adjustment is appropriate in patients with hepatic impairment. Remdesivir should only be used in patients with hepatic impairment if the potential benefit outweighs the potential risk. Liver function should be determined in all patients prior to starting remdesivir and should be monitored while receiving it as clinically appropriate.</p> <p>The SmPC recommends that remdesivir should not be initiated in patients with ALT $\geq 5x$ ULN at baseline. Remdesivir should be discontinued in patients who develop ALT $\geq 5x$ ULN (SmPC) or $>10x$ ULN (USPI) during treatment with remdesivir. Remdesivir should be discontinued in patients who develop ALT elevation accompanied by signs or symptoms of liver inflammation or increasing conjugated bilirubin, alkaline phosphatase, or international normalised ratio (INR).</p>
Ribavirin	Discontinue if progressive and clinically significant ALT rises, despite dose reduction, or accompanied by increased bilirubin.
Sotrovimab	No clinical trials have been conducted to evaluate the effects of hepatic impairment on the pharmacokinetics of sotrovimab. The impact of hepatic impairment on sotrovimab is unknown.
Tixagevimab/ Cilgavimab	No specific studies have been conducted to examine the effects of hepatic impairment on the pharmacokinetics of tixagevimab and cilgavimab. The effect of hepatic impairment on the pharmacokinetics of tixagevimab and cilgavimab is unknown.

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment

Charts updated 17 January 2022

Page 4 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.
No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

COVID-19 Host-directed Therapies (Licensed and Under Clinical Investigation)

Please refer to the additional notes on the following pages for further details and complete dose recommendations.

Drug	Hepatic Impairment			References
	Mild (Childs Pugh A)	Moderate (Childs Pugh B)	Severe (Childs Pugh C)	
Anakinra	100%	100%	100%, with caution	29, 30
Baricitinib	100%	100%	Not recommended	31, 32
Budesonide (inhaled)	Not studied. Consider 100% with close monitoring.	Not studied. Consider 100% with close monitoring.	Not studied. Consider 100% with close monitoring.	33, 34
Canakinumab	100%	100%	100%	35, 36
Colchicine	100% with close monitoring Contraindicated with P-gp inhibitors or strong CYP3A4 inhibitors.	100% with close monitoring Contraindicated with P-gp inhibitors or strong CYP3A4 inhibitors.	Contraindicated (SmPC) Consider dose reduction (USPI) Contraindicated with P-gp inhibitors or strong CYP3A4 inhibitors.	37, 38
Dexamethasone (low dose)	100%	100%	100%, with caution	39, 40
Hydrocortisone	Not studied. Consider dose reduction.	Not studied. Consider dose reduction.	Not studied. Consider dose reduction.	41
Infliximab	Not studied. No dosing recommendations can be made.	Not studied. No dosing recommendations can be made.	Not studied. No dosing recommendations can be made.	42, 43
Methylprednisolone	Not studied. Consider 100%.	Not studied. Consider 100%.	Not studied. Consider 100%, with caution	44, 45
Ruxolitinib	50% (dosing as per platelet count in product label)	50% (dosing as per platelet count in product label)	50% (dosing as per platelet count in product label)	46-48
Sarilumab	Not studied. Consider 100% under specialist supervision with frequent ALT monitoring.	Not studied. Consider 100% under specialist supervision with frequent ALT monitoring.	Not studied. Consider 100% under specialist supervision with frequent ALT monitoring.	49
Tocilizumab	Not studied. Consider 100% under specialist supervision with frequent ALT monitoring.	Not studied. Consider 100% under specialist supervision with frequent ALT monitoring.	Not studied. Consider 100% under specialist supervision with frequent ALT monitoring.	50, 51

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment

Charts updated 17 January 2022

Page 5 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.
No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

Notes - Host-directed Therapies (Licensed and Under Clinical Investigation)

Anakinra	The efficacy and safety in patients with AST/ALT ≥ 1.5 x ULN have not been evaluated.
Baricitinib	There was no clinically relevant effect on the pharmacokinetics of baricitinib in patients with mild or moderate hepatic impairment. The use of baricitinib has not been studied in patients with severe hepatic impairment.
Budesonide (inhaled)	Formal pharmacokinetic studies with inhaled budesonide have not been conducted in patients with hepatic impairment. However, since budesonide undergoes mainly hepatic metabolism, impairment of liver function may lead to accumulation of budesonide in the plasma. Patients with hepatic disease should be closely monitored.
Canakinumab	No formal pharmacokinetic studies have been performed in patients with hepatic impairment. Elimination of protein drugs such as canakinumab is thought to occur via proteolytic catabolism in different tissues. Although the liver is known to be a major organ of protein degradation, impaired hepatic function is not expected to be a limiting factor for elimination.
Colchicine	Use with caution in patients with mild/moderate hepatic impairment. Such patients should be carefully monitored for adverse effects of colchicine. Coadministration is contraindicated in the SmPC for patients with severe hepatic impairment, but the USPI recommends to consider dose reduction. Coadministration is contraindicated in patients with hepatic impairment who are taking a P-gp inhibitor or a strong CYP3A4 inhibitor.
Dexamethasone (low dose)	Particular care is required when considering the use of systemic corticosteroids in patients with liver failure and frequent patient monitoring is necessary. The elimination half-life is prolonged in severe liver disease.
Hydrocortisone	There may be an increased effect in patients with liver disease, and monitoring is advised. Reduced dosing may be considered.
Infliximab	Infliximab has not been studied in patients with hepatic impairment. No dose recommendations can be made.
Methylprednisolone	Particular care is required when considering the use of systemic corticosteroids in patients with liver failure or cirrhosis and frequent patient monitoring is necessary.
Ruxolitinib	Recommendations as for polycythaemia vera indication in product labels.
Sarilumab	Initiating treatment is not recommended in patients with ALT or AST >1.5 x ULN.
Tocilizumab	The European product label does not recommend treatment in patients with baseline ALT or AST >5 x ULN. The American product label does not recommend to initiate treatment in patients with elevated transaminases ALT or AST >1.5 x ULN.

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment

Charts updated 17 January 2022

Page 6 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.

No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

References

1. Zithromax (azithromycin) [Summary of Product Characteristics](#), Pfizer Ltd.
2. Zithromax (azithromycin) [Prescribing Information](#), Pfizer Inc.
3. Bamlanivimab and etesevimab [FDA Emergency Use Authorization](#), Lilly USA.
4. Ronapreve (casirivimab/imdevimab) [Summary of Product Characteristics](#), Roche Products Ltd.
5. REGEN-COV (casirivimab/imdevimab) [FDA Emergency Use Authorization](#), Regeneron Pharmaceuticals Inc.
6. Avloclor (chloroquine) tablets [Summary of Product Characteristics](#), Alliance Pharmaceuticals.
7. Aralen (chloroquine phosphate) tablets, [US Prescribing Information](#), Sanofi Aventis (discontinued).
8. Avigan (favipiravir) [Japanese Product Label](#), Toyama Chemical Co Ltd.
9. Pharmacokinetics of favipiravir in volunteers with hepatic impairment. [ClinicalTrials.gov Identifier: NCT01419457](#).
10. Plaquenil (hydroxychloroquine) [Summary of Product Characteristics](#), Zentiva.
11. Plaquenil (hydroxychloroquine) [US Prescribing Information](#), Concordia.
12. Rebif (interferon beta 1a) [Summary of Product Characteristics](#), Merck.
13. Rebif (interferon beta 1a) [US Prescribing Information](#), EMD Serono.
14. Betaferon (interferon beta 1b) [Summary of Product Characteristics](#), Bayer.
15. Betaseron (interferon beta 1b) [US Prescribing Information](#), Bayer HealthCare Pharmaceuticals Inc.
16. Soolantra (ivermectin) [Summary of Product Characteristics](#), Galderma (UK) Ltd.
17. Lagevrio (molnupiravir) [Summary of Product Characteristics](#), Merck Sharp and Dohme Ltd.
18. Yomesan (niclosamide) [Prescribing Information](#), Bayer (discontinued).
19. Paxlovid (nirmatrelvir/ritonavir) [Summary of Product Characteristics](#), Pfizer Ltd.
20. Paxlovid (nirmatrelvir/ritonavir) [FDA Emergency Use Authorization](#), Pfizer.
21. Alinia (nitazoxanide) [US Prescribing Information](#), Romark Pharmaceuticals.
22. Veklury (remdesivir) [Summary of Product Characteristics](#), Gilead Sciences Ltd.
23. Veklury (remdesivir) [US Prescribing Information](#), Gilead Sciences Inc.
24. Rebetol (ribavirin) [Summary of Product Characteristics](#), Merck Sharpe & Dohme Ltd.
25. Rebetol (ribavirin) [US Prescribing Information](#), Merck & Co Inc.
26. Xevudy (sotrovimab) [Summary of Product Characteristics](#), GlaxoSmithKline.
27. Sotrovimab [FDA Emergency Use Authorization](#), GlaxoSmithKline.
28. Evusheld (tixagevimab/cilgavimab) [FDA Emergency Use Authorization](#), AstraZeneca.

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment

Charts updated 17 January 2022

Page 7 of 7

Please check www.covid19-druginteractions.org for updates. Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.

No recommendation to use experimental therapy for COVID-19 is made. Data for many agents are limited or absent; therefore, risk-benefit assessment for any individual patient rests with prescribers.

29. Kineret (anakinra) [Summary of Product Characteristics](#), Swedish Orphan Biovitrum.
30. Kineret (anakinra) [US Prescribing Information](#), Swedish Orphan Biovitrum.
31. Olumiant (baricitinib) [Summary of Product Characteristics](#), Lilly.
32. Olumiant (baricitinib) [US Prescribing Information](#), Lilly USA.
33. Pulmicort (budesonide) [Summary of Product Characteristics](#), Astra Zeneca.
34. Pulmicort (budesonide) [US Prescribing Information](#), Astra Zeneca.
35. Ilaris (canakinumab) [US Prescribing Information](#), Novartis Pharmaceuticals.
36. Ilaris (canakinumab) [CHMP Assessment Report](#), European Medicines Agency.
37. Colchicine [Summary of Product Characteristics](#), Wockhardt UK Ltd.
38. Colcrys (colchicine) [US Prescribing Information](#), Takeda Pharmaceuticals America Inc.
39. Dexamethasone [Summary of Product Characteristics](#), Aspen.
40. Dexamethasone [Summary of Product Characteristics](#), Consilient Health Ltd.
41. Solu-Cortef (hydrocortisone) [Summary of Product Characteristics](#), Pfizer Ltd.
42. Remicade(infliximab) [Summary of Product Characteristics](#), Merck Sharp and Dohme Ltd.
43. Remicade (infliximab) [US Prescribing Information](#), Janssen Pharmaceuticals.
44. Medrone (methylprednisolone) [Summary of Product Characteristics](#), Pfizer Ltd.
45. Medrol (methylprednisolone) [US Prescribing Information](#), Pfizer.
46. Jakavi (ruxolitinib) [Summary of Product Characteristics](#), Novartis Pharmaceuticals Ltd.
47. Jakafi (ruxolitinib) [US Prescribing Information](#), Incyte Corporation.
48. Pharmacokinetics and pharmacodynamics of orally administered ruxolitinib (INCB018424 phosphate) in renal and hepatic impairment patients. Chen X, Shi JG, Emm T, *et al.* [Clin Pharmacol Drug Dev.](#) 2014;3(1):34-42.
49. Kevzara (sarilumab) [Summary of Product Characteristics](#), Sanofi Genzyme.
50. RoActemra (tocilizumab, for infusion) [Summary of Product Characteristics](#), Roche Products Ltd.
51. Actemra (tocilizumab) [US Prescribing Information](#), Genentech Inc.